

Duality for Knizhnik-Zamolodchikov and Dynamical Equations

V. Tarasov^{*} and A. Varchenko^{*}

^{*}*St. Petersburg Branch of Steklov Mathematical Institute
Fontanka 27, St. Petersburg 191011, Russia*

^{*}*Department of Mathematics, University of North Carolina
Chapel Hill, NC 27599, USA*

June 2001

Abstract. We consider the Knizhnik-Zamolodchikov (KZ) and dynamical equations, both differential and difference, in the context of the $(\mathfrak{gl}_k, \mathfrak{gl}_n)$ duality. We show that the KZ and dynamical equations naturally exchange under the duality.

Introduction

The Knizhnik-Zamolodchikov (KZ) equations is a holonomic system of differential equations for correlation functions in conformal field theory on the sphere. They play an important role in representation theory of affine Lie algebras and quantum groups. The difference analogue of the KZ equations is called the quantized Knizhnik-Zamolodchikov (q KZ) equations.

There are rational, trigonometric and elliptic versions of KZ and q KZ equations, depending on what kind of coefficient functions the equations have. In this paper we consider the rational and trigonometric versions.

In [FMTV] we discovered another holonomic system of differential equations with rational coefficients, which is compatible with an extension of the rational KZ equations by an element of the Cartan subalgebra. We call the new equations the rational *dynamical differential* (DD) equations. The difference analogue of the DD equations was constructed in [TV1] and called the rational *difference dynamical* (q DD) equations. The rational q DD equations are compatible with the trigonometric KZ equations.

Around 1995 for a simple Lie algebra $\mathfrak{g}$ De Concini and Procesi introduced a $U(\mathfrak{g})$ valued connection on the set of regular elements of the Cartan subalgebra of $\mathfrak{g}$ [CP]. They conjectured that the monodromy of the connection is described in terms of the corresponding quantum Weyl group. In the recent work of Toledano-Laredo this conjecture is proved for $\mathfrak{g} = \mathfrak{sl}_n$ [TL].

When the work [FMTV] had been written, P. Etingof told us about the De Concini–Procesi connection and indicated that their connection is a special case of our DD equations.

^{*}Supported in part by RFFI grant 99–01–00101 and INTAS grant 99–01705

E-mail: vt@pdmi.ras.ru

^{*}Supported in part by NSF grant DMS–9801582

E-mail: anv@email.unc.edu

The $(\mathfrak{gl}_k, \mathfrak{gl}_n)$ duality plays an important role in the representation theory and the classical invariant theory, see [Zh1], [H]. In [TL] Toledano-Laredo discovered that with respect to this duality the De Concini–Procesi connection is dual to the rational differential KZ equations and used this fact to compute the monodromy of the De Concini–Procesi connection in terms of the quantum Weyl group. Our paper is inspired by the result of [TL].

In this paper we show that under the $(\mathfrak{gl}_k, \mathfrak{gl}_n)$ duality the KZ and qKZ equations for the Lie algebra $\mathfrak{gl}_k$ exchange with the DD and qDD equations for the Lie algebra $\mathfrak{gl}_n$, while the DD and qDD equations for $\mathfrak{gl}_k$ exchange with the KZ and qKZ equations for $\mathfrak{gl}_n$.

The duality between the KZ and DD equations in the rational differential case is essentially the “quantum” version of the duality for isomonodromic deformation systems [Ha1]. We thank J. Harnad for pointing out this observation to us. The relation of the differential KZ equations and the isomonodromic deformation systems is described in [R], [Ha2].

An interplay between extremal projectors and R -matrices, relevant to the duality studied in our paper, has been observed in [ST].

We consider three cases in the paper: the rational differential, trigonometric differential, and rational difference cases. To consider the trigonometric difference case, the duality for the trigonometric qKZ equations and the trigonometric qDD equations, one has to employ the q -analogue of the $(\mathfrak{gl}_k, \mathfrak{gl}_n)$ duality: the $(U_q(\mathfrak{gl}_k), U_q(\mathfrak{gl}_n))$ duality. This will be done elsewhere. The $(U_q(\mathfrak{gl}_k), U_q(\mathfrak{gl}_n))$ duality has been described in [B], [TL].

The paper is organized as follows. After introducing basic notation we subsequently describe the differential KZ and DD equations, and the rational difference qKZ and qDD equations, for the Lie algebra $\mathfrak{gl}_k$. This is done in Sections 2–4. The main results are Theorems 5.8 and 5.9 in Section 5.

The first author thanks Maxim Nazarov for helpful discussions.

1. Basic notation

Let n be a nonnegative integer. A *partition* $\lambda = (\lambda_1, \lambda_2, \dots)$ with at most k parts is an infinite nonincreasing sequence of nonnegative integers such that $\lambda_{k+1} = 0$. Denote by $\mathcal{P}_k$ the set of partitions with at most k parts and by $\mathcal{P}$ the set of all partitions. We often make use of the embedding $\mathcal{P}_k \rightarrow \mathbb{C}^k$ given by truncating the zero tail of a partition: $(\lambda_1, \dots, \lambda_k, 0, 0, \dots) \mapsto (\lambda_1, \dots, \lambda_k)$. Since obviously $\mathcal{P}_m \subset \mathcal{P}_k$ for $m \leq k$, in fact, we have a collection of embeddings $\mathcal{P}_m \rightarrow \mathbb{C}^k$ for any $m \leq k$. What particular embedding is used will be clear from the context.

Let e_{ab} , $a, b = 1, \dots, k$, be the standard basis of the Lie algebra $\mathfrak{gl}_k$: $[e_{ab}, e_{cd}] = \delta_{bc}e_{ad} - \delta_{ad}e_{cb}$. We take the Cartan subalgebra $\mathfrak{h} \subset \mathfrak{gl}_k$ spanned by $e_{11}, \dots, e_{kk}$, and the nilpotent subalgebras $\mathfrak{n}_+$ and $\mathfrak{n}_-$ spanned by the elements e_{ab} for $a < b$ and $a > b$, respectively. We have the standard Gauss decomposition $\mathfrak{gl}_k = \mathfrak{n}_+ \oplus \mathfrak{h} \oplus \mathfrak{n}_-$.

Let $\varepsilon_1, \dots, \varepsilon_k$ be the basis of $\mathfrak{h}^*$ dual to $e_{11}, \dots, e_{kk}$: $\langle \varepsilon_a, e_{bb} \rangle = \delta_{ab}$. We identify $\mathfrak{h}^*$ with $\mathbb{C}^k$ mapping $\lambda_1 \varepsilon_1 + \dots + \lambda_k \varepsilon_k$ to $(\lambda_1, \dots, \lambda_k)$. The root vectors of $\mathfrak{gl}_k$ are e_{ab} for $a \neq b$, the corresponding root being equal to $\alpha_{ab} = \varepsilon_a - \varepsilon_b$. The roots α_{ab} for $a < b$ are positive. The simple roots are $\alpha_1, \dots, \alpha_{k-1}$: $\alpha_a = \varepsilon_a - \varepsilon_{a+1}$. Denote by ρ a half-sum of positive roots.

We choose the standard invariant bilinear form $(,)$ on $\mathfrak{gl}_k$: $(e_{ab}, e_{cd}) = \delta_{ad}\delta_{bc}$. It defines an isomorphism $\mathfrak{h} \rightarrow \mathfrak{h}^*$. The induced bilinear form on $\mathfrak{h}^*$ is $(\varepsilon_a, \varepsilon_b) = \delta_{ab}$.

The element $h = e_{11} + \dots + e_{kk} \in \mathfrak{h}$ is central in $\mathfrak{gl}_k$. We identify the Lie algebra $\mathfrak{sl}_k$ with the Lie subalgebra of $\mathfrak{gl}_k$ orthogonal to h . Then $\mathfrak{gl}_k = \mathfrak{sl}_k \oplus \mathbb{C}h$.

For a $\mathfrak{gl}_k$ -module W and a weight $\lambda \in \mathfrak{h}^*$ let $W[\lambda]$ be the weight subspace of W of weight λ .

For any $\lambda \in \mathcal{P}_k$ we denote by V_λ the irreducible $\mathfrak{gl}_k$ -module with highest weight λ . By abuse of notation, for any $l \in \mathbb{Z}_{\geq 0}$ we write V_l instead of $V_{(l,0,\dots,0)}$. Thus, $V_0 = \mathbb{C}$ is the trivial $\mathfrak{gl}_k$ -module, $V_1 = \mathbb{C}^k$ with the natural action of $\mathfrak{gl}_k$, and V_l is the l -th symmetric power of V_1 .

The element $I = \sum_{a,b=1}^k e_{ab}e_{ba}$ is central in $U(\mathfrak{gl}_k)$. In the irreducible $\mathfrak{gl}_k$ -module V_λ it acts as multiplication by $(\lambda, \lambda + 2\rho)$.

Define a $\mathfrak{gl}_k$ -action on the polynomial ring $\mathbb{C}[x_1, \dots, x_k]$ by differential operators: $e_{ab} \mapsto x_a \partial_b$, where $\partial_b = \partial/\partial x_b$, and denote the obtained $\mathfrak{gl}_k$ -module by $V_\bullet$. Then

$$(1.1) \quad V_\bullet = \bigoplus_{l=0}^{\infty} V_l,$$

the submodule V_l being spanned by homogeneous polynomials of degree l . The highest weight vector of the submodule V_l is x_1^l .

2. Knizhnik-Zamolodchikov and differential dynamical operators

For any $g \in U(\mathfrak{gl}_k)$ set $g_{(i)} = \text{id} \otimes \dots \otimes g \otimes \dots \otimes \text{id} \in (U(\mathfrak{gl}_k))^{\otimes n}$. We consider $U(\mathfrak{gl}_k)$ as a subalgebra of $(U(\mathfrak{gl}_k))^{\otimes n}$, the embedding $U(\mathfrak{gl}_k) \hookrightarrow (U(\mathfrak{gl}_k))^{\otimes n}$ being given by the n -fold coproduct, that is, $x \mapsto x_{(1)} + \dots + x_{(n)}$ for any $x \in \mathfrak{gl}_k$.

Let $\Omega = \sum_{a,b=1}^k e_{ab} \otimes e_{ba}$ be the Casimir tensor, and let $\Omega^0 = \frac{1}{2} \sum_{a=1}^k e_{aa} \otimes e_{aa}$,

$$\Omega^+ = \Omega^0 + \sum_{1 \leq a < b \leq k} e_{ab} \otimes e_{ba}, \quad \Omega^- = \Omega^0 + \sum_{1 \leq a < b \leq k} e_{ba} \otimes e_{ab},$$

so that $\Omega = \Omega^+ + \Omega^-$.

Fix a nonzero complex number κ . Consider differential operators $\nabla_{z_1}, \dots, \nabla_{z_n}$ and $\hat{\nabla}_{z_1}, \dots, \hat{\nabla}_{z_n}$ with coefficients in $(U(\mathfrak{gl}_k))^{\otimes n}$ depending on complex variables $z_1, \dots, z_n$ and $\lambda_1, \dots, \lambda_k$:

$$\begin{aligned} \nabla_{z_i}(z; \lambda) &= \kappa \frac{\partial}{\partial z_i} - \sum_{a=1}^k \lambda_a (e_{aa})_{(i)} - \sum_{\substack{j=1 \\ j \neq i}}^n \frac{\Omega_{(ij)}}{z_i - z_j}, \\ \hat{\nabla}_{z_i}(z; \lambda) &= \kappa z_i \frac{\partial}{\partial z_i} - \sum_{a=1}^k \left(\lambda_a - \frac{e_{aa}}{2} \right) (e_{aa})_{(i)} - \sum_{\substack{j=1 \\ j \neq i}}^n \frac{z_i \Omega_{(ij)}^+ + z_j \Omega_{(ij)}^-}{z_i - z_j}. \end{aligned}$$

The differential operators $\nabla_{z_1}, \dots, \nabla_{z_n}$ (resp. $\widehat{\nabla}_{z_1}, \dots, \widehat{\nabla}_{z_n}$) are called the *rational* (resp. *trigonometric*) *Knizhnik-Zamolodchikov (KZ) operators*. The following statements are well known.

Theorem 2.1. *The operators $\nabla_{z_1}, \dots, \nabla_{z_n}$ pairwise commute.*

Theorem 2.2. *The operators $\widehat{\nabla}_{z_1}, \dots, \widehat{\nabla}_{z_n}$ pairwise commute.*

The *rational KZ equations* is a system of differential equations

$$\nabla_{z_i} u = 0, \quad i = 1, \dots, n,$$

for a function $u(z_1, \dots, z_n; \lambda_1, \dots, \lambda_k)$ taking values in an n -fold tensor product of $\mathfrak{gl}_k$ -modules. The *trigonometric KZ equations* is a system of differential equations

$$\widehat{\nabla}_{z_i} u = 0, \quad i = 1, \dots, n.$$

Introduce differential operators $D_{\lambda_1}, \dots, D_{\lambda_k}$ and $\widehat{D}_{\lambda_1}, \dots, \widehat{D}_{\lambda_k}$ with coefficients in $(U(\mathfrak{gl}_k))^{\otimes n}$ depending on complex variables $z_1, \dots, z_n$ and $\lambda_1, \dots, \lambda_k$:

$$\begin{aligned} D_{\lambda_a}(z; \lambda) &= \kappa \frac{\partial}{\partial \lambda_a} - \sum_{i=1}^n z_i (e_{aa})_{(i)} - \sum_{\substack{b=1 \\ b \neq a}}^k \frac{e_{ab} e_{ba} - e_{aa}}{\lambda_a - \lambda_b}. \\ \widehat{D}_{\lambda_a}(z; \lambda) &= \kappa \lambda_a \frac{\partial}{\partial \lambda_a} + \frac{e_{aa}^2}{2} - \sum_{i=1}^n z_i (e_{aa})_{(i)} - \\ &\quad - \sum_{b=1}^k \sum_{1 \leq i < j \leq n} (e_{ab})_{(i)} (e_{ba})_{(j)} - \sum_{\substack{b=1 \\ b \neq a}}^k \frac{\lambda_b}{\lambda_a - \lambda_b} (e_{ab} e_{ba} - e_{aa}). \end{aligned}$$

Recall that $e_{ab} = \sum_{i=1}^n (e_{ab})_{(i)}$. We call the differential operators $D_{\lambda_1}, \dots, D_{\lambda_k}$ (resp. $\widehat{D}_{\lambda_1}, \dots, \widehat{D}_{\lambda_k}$) the *rational* (resp. *trigonometric*) *differential dynamical (DD) operators*.

Theorem 2.3. *The operators $\nabla_{z_1}, \dots, \nabla_{z_n}, D_{\lambda_1}, \dots, D_{\lambda_k}$ pairwise commute.*

The statement follows from the same result for the $\mathfrak{sl}_k$ case in [FMTV].

Theorem 2.4. [TV2] *The operators $\widehat{D}_{\lambda_1}, \dots, \widehat{D}_{\lambda_k}$ pairwise commute.*

Later we will formulate analogues of Theorem 2.3 for the trigonometric KZ operators and for the trigonometric DD operators. They involve difference dynamical operators and difference (quantized) Knizhnik-Zamolodchikov operators.

The *rational DD equations* is a system of differential equations

$$D_{\lambda_a} u = 0, \quad a = 1, \dots, k,$$

for a function $u(z_1, \dots, z_n; \lambda_1, \dots, \lambda_k)$ taking values in an n -fold tensor product of $\mathfrak{gl}_k$ -modules. The *trigonometric DD equations* is a system of differential equations

$$\widehat{D}_{\lambda_a} u = 0, \quad a = 1, \dots, k.$$

3. Rational difference dynamical operators

For any $a, b = 1, \dots, k$, $a \neq b$, introduce a series $B_{ab}(t)$ depending on a complex variable t :

$$B_{ab}(t) = 1 + \sum_{s=1}^{\infty} e_{ba}^s e_{ab}^s \prod_{j=1}^s \frac{1}{j(t - e_{aa} + e_{bb} - j)}.$$

The series has a well-defined action in any finite-dimensional $\mathfrak{gl}_k$ -module W , giving an $\text{End}(W)$ -valued rational function of t . These series have zero weight:

$$[B_{ab}(t), x] = 0 \quad \text{for any } x \in \mathfrak{h},$$

satisfy the inversion relation

$$B_{ab}(t) B_{ba}(-t) = 1 - \frac{e_{aa} - e_{bb}}{t},$$

and the braid relation

$$B_{ab}(t) B_{ac}(t + s) B_{bc}(s) = B_{bc}(s) B_{ac}(t + s) B_{ab}(t).$$

Remark. In notation of [TV1] the series $B_{ab}(t)$ equals $p(t-1; e_{aa} - e_{bb}, e_{ab}, e_{ba})$. The properties of $B_{ab}(t)$ follow from the properties of functions $B_w(\lambda)$ in the $\mathfrak{sl}_k$ case, see [TV1, Section 2.6].

Remark. The series $B_{ab}(t)$ first appeared in the definition of the extremal cocycle in [Zh2].

Consider the products $X_1, \dots, X_k$ depending on complex variables $z_1, \dots, z_n$ and $\lambda_1, \dots, \lambda_k$:

$$(3.1) \quad X_a(z; \lambda) = (B_{ak}(\lambda_{ak}) \dots B_{a,a+1}(\lambda_{a,a+1}))^{-1} \times \\ \times \prod_{i=1}^n (z_i^{-e_{aa}})_{(i)} B_{1a}(\lambda_{1a} - \kappa) \dots B_{a-1,a}(\lambda_{a-1,a} - \kappa),$$

acting in any n -fold tensor product $W_1 \otimes \dots \otimes W_n$ of finite-dimensional $\mathfrak{gl}_k$ -modules. Here $\lambda_{bc} = \lambda_b - \lambda_c$.

Denote by T_u a difference operator acting on a function $f(u)$ by

$$(T_u f)(u) = f(u + \kappa).$$

Introduce difference operators $Q_{\lambda_1}, \dots, Q_{\lambda_k}$:

$$Q_{\lambda_a}(z; \lambda) = X_a(z; \lambda) T_{\lambda_a}.$$

We call these operators the (*rational*) *difference dynamical* (qDD) operators.

Theorem 3.1. *The operators $\widehat{\nabla}_{z_1}, \dots, \widehat{\nabla}_{z_n}$, $Q_{\lambda_1}, \dots, Q_{\lambda_k}$ pairwise commute.*

The statement follows from the same result for the $\mathfrak{sl}_k$ case in [TV1].

In more conventional form the equalities $[\widehat{\nabla}_{z_i}, Q_{\lambda_a}] = 0$ and $[Q_{\lambda_a}, Q_{\lambda_b}] = 0$ respectively look like:

$$\widehat{\nabla}_{z_i}(z; \lambda) X_a(z; \lambda) = X_a(z; \lambda) \widehat{\nabla}_{z_i}(z; \lambda_1, \dots, \lambda_a + \kappa, \dots, \lambda_k),$$

$$X_a(z; \lambda) X_b(z; \lambda_1, \dots, \lambda_a + \kappa, \dots, \lambda_k) = X_b(z; \lambda) X_a(z; \lambda_1, \dots, \lambda_b + \kappa, \dots, \lambda_k).$$

The *rational qDD equations* is a system of difference equations

$$Q_{\lambda_a} u = u, \quad a = 1, \dots, k,$$

for a function $u(z_1, \dots, z_n; \lambda_1, \dots, \lambda_k)$ taking values in an n -fold tensor product of $\mathfrak{gl}_k$ -modules.

4. Rational difference Knizhnik-Zamolodchikov operators

Consider the Yangian $Y(\mathfrak{gl}_k)$. The algebra $U(\mathfrak{gl}_k)$ is embedded in $Y(\mathfrak{gl}_k)$ as a Hopf subalgebra, and we identify $U(\mathfrak{gl}_k)$ with the image of this embedding.

There is an algebra homomorphism $ev : Y(\mathfrak{gl}_k) \rightarrow U(\mathfrak{gl}_k)$, called the *evaluation homomorphism*, which is identical on the subalgebra $U(\mathfrak{gl}_k) \subset Y(\mathfrak{gl}_k)$. The evaluation homomorphism is not a homomorphism of Hopf algebras.

The Yangian $Y(\mathfrak{gl}_k)$ has a distinguished one-parametric family of automorphisms ρ_u depending on a complex parameter u . For any $\mathfrak{gl}_k$ -module W we denote by $W(u)$ the pullback of W through the homomorphism $ev \circ \rho_u$. Yangian modules of this form are called *evaluation modules*.

For any finite-dimensional irreducible $\mathfrak{gl}_k$ -modules V, W the tensor products $V(t) \otimes W(u)$ and $W(u) \otimes V(t)$ are isomorphic irreducible $Y(\mathfrak{gl}_k)$ -modules, provided $t - u \notin \mathbb{Z}$. The intertwiner can be taken of the form $P_{VW} R_{VW}(t - u)$ where $P_{VW} : V \otimes W \rightarrow W \otimes V$ is the flip map: $P_{VW} : v \otimes w \mapsto w \otimes v$, and $R_{VW}(t)$ is a rational $\text{End}(V \otimes W)$ valued function, called the *rational R-matrix* for the tensor product $V \otimes W$.

We describe below the R -matrix $R_{VW}(t)$ in terms of the $\mathfrak{gl}_k$ actions in V and W . Let v and w be the respective highest weight vectors. The R -matrix $R_{VW}(t)$ is uniquely determined by the $\mathfrak{gl}_k$ -invariance,

$$(4.1) \quad [R_{VW}(t), g \otimes 1 + 1 \otimes g] = 0 \quad \text{for any } g \in \mathfrak{gl}_k,$$

the commutation relations

$$(4.2) \quad R_{VW}(t) (te_{ab} \otimes 1 + \sum_{c=1}^k e_{ac} \otimes e_{cb}) = (te_{ab} \otimes 1 + \sum_{c=1}^k e_{cb} \otimes e_{ac}) R_{VW}(t),$$

and the normalization condition

$$R_{VW}(t) v \otimes w = v \otimes w.$$

The introduced R -matrices obey the inversion relation

$$(4.3) \quad R_{VW}(t) R_{WV}^{(21)}(-t) = 1$$

where $R_{WV}^{(21)} = P_{WV} R_{WV} P_{VW}$, and the Yang-Baxter equation

$$(4.4) \quad R_{UV}(t) R_{UW}(t+u) R_{VW}(u) = R_{VW}(u) R_{UW}(t+u) R_{UV}(t).$$

The aforementioned facts on the Yangian $Y(\mathfrak{gl}_k)$ are well known; for example, see [MNO].

Consider the $\mathfrak{gl}_k$ -module $V_\bullet$, and let $v_l = x_1^l$ be the highest weight vector of the irreducible component $V_l \subset V_\bullet$. We define the R -matrix $R_{V_\bullet V_\bullet}(t)$ as a direct sum of the R -matrices $R_{V_l V_m}(t)$:

$$R_{V_\bullet V_\bullet}(t) v \otimes v' = R_{V_l V_m}(t) v \otimes v' \quad \text{for any } v \otimes v' \in V_l \otimes V_m,$$

the R -matrices $R_{V_l V_m}(t)$ being normalized by $R_{V_l V_m}(t) v_l \otimes v_m = v_l \otimes v_m$. It is clear that $R_{V_\bullet V_\bullet}(t)$ obeys both the inversion relation and the Yang-Baxter equation.

Consider the products $K_1, \dots, K_n$ depending on complex variables $z_1, \dots, z_n$ and $\lambda_1, \dots, \lambda_k$:

$$(4.5) \quad K_i(z; \lambda) = (R_{in}(z_{in}) \dots R_{i,i+1}(z_{i,i+1}))^{-1} \times \\ \times \prod_{a=1}^k (\lambda_a^{-e_{aa}})_{(i)} R_{1i}(z_{1i} - \kappa) \dots R_{i-1,i}(z_{i-1,i} - \kappa),$$

acting in an n -fold tensor product $W_1 \otimes \dots \otimes W_n$ of $\mathfrak{gl}_k$ -modules. Here $z_{ij} = z_i - z_j$, and $R_{ij}(t) = (R_{W_i W_j}(t))_{(ij)}$.

Introduce difference operators $Z_{z_1}, \dots, Z_{z_n}$:

$$Z_{z_i}(z; \lambda) = K_i(z; \lambda) T_{z_i}.$$

We call these operators the (*rational*) *quantized Knizhnik-Zamolodchikov* (qKZ) operators.

Theorem 4.1. [FR] *The operators $Z_{z_1}, \dots, Z_{z_n}$ pairwise commute.*

Theorem 4.2. [TV2] *The operators $Z_{z_1}, \dots, Z_{z_n}$, $\widehat{D}_{\lambda_1}, \dots, \widehat{D}_{\lambda_k}$ pairwise commute.*

The last theorem extends Theorems 2.4 and 4.1.

In more conventional form the equalities $[Z_{z_i}, Z_{z_j}] = 0$ and $[Z_{z_i}, \widehat{D}_{\lambda_a}] = 0$ respectively look like:

$$K_i(z; \lambda) K_j(z_1, \dots, z_i + \kappa, \dots, z_n; \lambda) = K_j(z; \lambda) K_i(z_1, \dots, z_j + \kappa, \dots, z_n; \lambda), \\ \widehat{D}_{\lambda_a}(z; \lambda) K_i(z; \lambda) = K_i(z; \lambda) \widehat{D}_{\lambda_a}(z_1, \dots, z_i + \kappa, \dots, z_n, \lambda).$$

The *rational qKZ equations* is a system of difference equations

$$Z_{z_i} u = u, \quad i = 1, \dots, n,$$

for a function $u(z_1, \dots, z_n; \lambda_1, \dots, \lambda_k)$ taking values in an n -fold tensor product of $\mathfrak{gl}_k$ -modules.

5. $(\mathfrak{gl}_k, \mathfrak{gl}_n)$ duality

In what follows we are going to consider the Lie algebras $\mathfrak{gl}_k$ and $\mathfrak{gl}_n$ simultaneously. In order to distinguish generators, modules, etc., we will indicate the dependence on k and n explicitly, for example, $e_{ab}^{(k)}$, $V_\lambda^{(n)}$.

Consider the polynomial ring $P_{kn} = \mathbb{C}[x_{11}, \dots, x_{k1}, \dots, x_{1n}, \dots, x_{kn}]$ of kn variables. We define a $\mathfrak{gl}_k$ -action on P_{kn} by

$$(5.1) \quad e_{ab}^{(k)} \mapsto \sum_{i=1}^n x_{ai} \partial_{bi},$$

where $\partial_{bi} = \partial / \partial x_{bi}$, and we define a $\mathfrak{gl}_n$ -action on P_{kn} by

$$(5.2) \quad e_{ij}^{(n)} \mapsto \sum_{a=1}^k x_{ai} \partial_{aj}.$$

There are two natural isomorphisms of vector spaces: $(\mathbb{C}[x_1, \dots, x_k])^{\otimes n} \rightarrow P_{kn}$,

$$(5.3) \quad (p_1 \otimes \dots \otimes p_n)(x_{11}, \dots, x_{kn}) = \prod_{i=1}^n p_i(x_{1i}, \dots, x_{ki}),$$

and $(\mathbb{C}[x_1, \dots, x_n])^{\otimes k} \rightarrow P_{kn}$,

$$(5.4) \quad (p_1 \otimes \dots \otimes p_k)(x_{11}, \dots, x_{kn}) = \prod_{a=1}^k p_a(x_{a1}, \dots, x_{an}).$$

We have a simple proposition.

Proposition 5.1. *The module P_{kn} is isomorphic to $(V_\bullet^{(k)})^{\otimes n}$ as a $\mathfrak{gl}_k$ -module by (5.3), and it is isomorphic to $(V_\bullet^{(n)})^{\otimes k}$ as a $\mathfrak{gl}_n$ -module by (5.4).*

It is easy to see that the actions (5.1) and (5.2) commute with each other, thus making P_{kn} into a module over the direct sum $\mathfrak{gl}_k \oplus \mathfrak{gl}_n$. The following theorem is well known, e.g. see [H].

Theorem 5.2. *The $\mathfrak{gl}_k \oplus \mathfrak{gl}_n$ module P_{kn} has the decomposition*

$$P_{kn} = \bigoplus_{\lambda \in \mathcal{P}_{\min(k,n)}} V_\lambda^{(k)} \otimes V_\lambda^{(n)}.$$

Fix vectors $\lambda = (l_1, \dots, l_n) \in \mathbb{Z}_{\geq 0}^n$ and $\mu = (m_1, \dots, m_k) \in \mathbb{Z}_{\geq 0}^k$ such that $\sum_{i=1}^n l_i = \sum_{a=1}^k m_a$. Let

$$\mathcal{Z}_{kn}[\lambda, \mu] = \left\{ (d_{ai})_{\substack{a=1, \dots, k \\ i=1, \dots, n}} \in \mathbb{Z}_{\geq 0}^{kn} \mid \sum_{a=1}^k d_{ai} = l_i, \quad \sum_{i=1}^n d_{ai} = m_a \right\}.$$

Denote by $P_{kn}[\lambda, \mu] \subset P_{kn}$ the span of all monomials $\prod_{a=1}^k \prod_{i=1}^n x_{ai}^{d_{ai}}$ such that $(d_{ai}) \in \mathcal{Z}_{kn}[\lambda, \mu]$. The next statement easily follows from Proposition 5.1, and formulae (1.1), (5.1) and (5.2).

Proposition 5.3. *The isomorphisms (5.3) and (5.4) induce the isomorphisms of the weight subspaces $(V_{l_1}^{(k)} \otimes \dots \otimes V_{l_n}^{(k)})[m_1, \dots, m_k]$ and $(V_{m_1}^{(n)} \otimes \dots \otimes V_{m_k}^{(n)})[l_1, \dots, l_n]$, and the space $P_{kn}[\lambda, \mu]$.*

In what follows we will need another description of the above mentioned isomorphisms. It is given below.

For an indeterminate y and a nonnegative integer s we define the *divided powers* $y^{[s]}$ by the rule: $y^{[0]} = 1$ and $y^{[s]} = y^s/s!$ for $s > 0$.

Let $v_{l_i}^{(k)}, v_{m_j}^{(n)}$ be highest weight vectors of the respective modules $V_{l_i}^{(k)}, V_{m_j}^{(n)}$.

Lemma 5.4. *A basis of the weight subspace $(V_{l_1}^{(k)} \otimes \dots \otimes V_{l_n}^{(k)})[m_1, \dots, m_k]$ is given by vectors*

$$v_{\mathbf{d}}^{(k)} = \prod_{a=2}^k (e_{a1}^{(k)})^{[d_{a1}]} v_{l_1}^{(k)} \otimes \dots \otimes \prod_{a=2}^k (e_{a1}^{(k)})^{[d_{an}]} v_{l_n}^{(k)}, \quad \mathbf{d} = (d_{ai}) \in \mathcal{Z}_{kn}[\lambda, \mu].$$

Lemma 5.5. *A basis of the weight subspace $(V_{m_1}^{(n)} \otimes \dots \otimes V_{m_k}^{(n)})[l_1, \dots, l_n]$ is given by vectors*

$$v_{\mathbf{d}}^{(n)} = \prod_{i=2}^n (e_{i1}^{(n)})^{[d_{1i}]} v_{m_1}^{(n)} \otimes \dots \otimes \prod_{i=2}^n (e_{i1}^{(n)})^{[d_{ki}]} v_{m_k}^{(n)}, \quad \mathbf{d} = (d_{ai}) \in \mathcal{Z}_{kn}[\lambda, \mu].$$

For any $\mathbf{d} \in \mathcal{Z}_{kn}[\lambda, \mu]$ set $x^{[\mathbf{d}]} = \prod_{a=1}^k \prod_{i=1}^n x_{ai}^{[d_{ai}]}$. Define isomorphisms of vector spaces:

$$(5.5) \quad (V_{l_1}^{(k)} \otimes \dots \otimes V_{l_n}^{(k)})[m_1, \dots, m_k] \rightarrow P_{kn}[\lambda, \mu], \quad v_{\mathbf{d}}^{(k)} \mapsto x^{[\mathbf{d}]},$$

$$(5.6) \quad (V_{m_1}^{(n)} \otimes \dots \otimes V_{m_k}^{(n)})[l_1, \dots, l_n] \rightarrow P_{kn}[\lambda, \mu], \quad v_{\mathbf{d}}^{(n)} \mapsto x^{[\mathbf{d}]},$$

and extend them by linearity to isomorphisms of vector spaces $(V_{\bullet}^{(k)})^{\otimes n} \rightarrow P_{kn}$ and $(V_{\bullet}^{(n)})^{\otimes k} \rightarrow P_{kn}$.

Proposition 5.6. *The maps (5.5) and (5.6) coincide with the maps (5.3) and (5.4), respectively.*

The proof is straightforward.

Consider the action of KZ , qKZ , DD and qDD operators for the Lie algebras $\mathfrak{gl}_k$ and $\mathfrak{gl}_n$ on P_{kn} -valued functions of $z_1, \dots, z_n$ and $\lambda_1, \dots, \lambda_k$, treating the space P_{kn} as a tensor product $(V_{\bullet}^{(k)})^{\otimes n}$ of $\mathfrak{gl}_k$ -modules, and as a tensor product $(V_{\bullet}^{(n)})^{\otimes k}$ of $\mathfrak{gl}_n$ -modules. We will write $F \simeq G$ if the operators F and G act on the P_{kn} -valued functions in the same way.

Lemma 5.7. *We have $I^{(k)} - k \sum_{a=1}^k e_{aa}^{(k)} \simeq I^{(n)} - n \sum_{i=1}^n e_{ii}^{(n)}$.*

The proof is straightforward.

Set

$$(5.7) \quad C_{ab}^{(k)}(t) = \frac{\Gamma(t+1) \Gamma(t - e_{aa}^{(k)} + e_{bb}^{(k)})}{\Gamma(t - e_{aa}^{(k)}) \Gamma(t + e_{bb}^{(k)} + 1)}, \quad C_{ij}^{(n)}(t) = \frac{\Gamma(t+1) \Gamma(t - e_{ii}^{(n)} + e_{jj}^{(n)})}{\Gamma(t - e_{ii}^{(n)}) \Gamma(t + e_{jj}^{(n)} + 1)}.$$

Theorem 5.8. For any $i = 1, \dots, n$ and $a = 1, \dots, k$ we have

$$(5.8) \quad \nabla_{z_i}^{(k)}(z; \lambda) \simeq D_{z_i}^{(n)}(\lambda; z), \quad D_{\lambda_a}^{(k)}(z; \lambda) \simeq \nabla_{\lambda_a}^{(n)}(\lambda; z),$$

$$(5.9) \quad \widehat{\nabla}_{z_i}^{(k)}(z; \lambda) \simeq \widehat{D}_{z_i}^{(n)}(\lambda; z), \quad \widehat{D}_{\lambda_a}^{(k)}(z; \lambda) \simeq \widehat{\nabla}_{\lambda_a}^{(n)}(\lambda; z),$$

$$(5.10) \quad Z_{z_i}^{(k)}(z; \lambda) \simeq N_i^{(n)}(z) Q_{z_i}^{(n)}(\lambda; z), \quad N_a^{(k)}(\lambda) Q_{\lambda_a}^{(k)}(z; \lambda) \simeq Z_{\lambda_a}^{(n)}(\lambda; z).$$

Here

$$N_i^{(n)}(z) = \prod_{1 \leq j < i} C_{ji}^{(n)}(z_{ji} - \kappa) \prod_{i < j \leq n} (C_{ij}^{(n)}(z_{ij}))^{-1}$$

and

$$N_a^{(k)}(\lambda) = \prod_{1 \leq b < a} C_{ba}^{(k)}(\lambda_{ba} - \kappa) \prod_{a < b \leq k} (C_{ab}^{(k)}(\lambda_{ab}))^{-1}.$$

Proof. Equalities (5.8) and (5.9) for differential operators are verified in a straightforward way. Equalities (5.10) for difference operators follow from Theorem 5.9, and formulae (3.1) and (4.5). $\square$

Remark. The first of equalities (5.8) at $\lambda = 0$ (respectively, the second one at $z = 0$) has been discovered in [TL].

Theorem 5.9. For any $a, b = 1, \dots, k$, $a \neq b$, and any $i, j = 1, \dots, n$, $i \neq j$, we have

$$(5.11) \quad B_{ab}^{(k)}(t) C_{ab}^{(k)}(t) \simeq R_{ab}^{(n)}(t), \quad R_{ij}^{(k)}(t) \simeq B_{ij}^{(n)}(t) C_{ij}^{(n)}(t).$$

Remark. Cf. the observation made after Theorem 6.6 in [FV].

Proof of Theorem 5.9. The Lie algebras $\mathfrak{gl}_k$ and $\mathfrak{gl}_n$ appear in our consideration on equal footing. So, it suffices to prove only the first formula in (5.11). Moreover, since both the action of $B_{ab}^{(k)}(t) C_{ab}^{(k)}(t)$ on P_{kn} and that of $R_{ab}^{(n)}(t)$ involves only the variables $x_{a1}, \dots, x_{an}$, $x_{b1}, \dots, x_{bn}$, it is enough to prove the claim for $k = 2$.

Let $\lambda = (l_1, \dots, l_n)$. Set $|\lambda| = l_1 + \dots + l_n$. Consider the decomposition of the $\mathfrak{gl}_2$ -module $V_{l_1}^{(2)} \otimes \dots \otimes V_{l_n}^{(2)}$ into a direct sum of irreducibles:

$$V_{l_1}^{(2)} \otimes \dots \otimes V_{l_n}^{(2)} = \bigoplus_{0 \leq m \leq |\lambda|/2} V_{(|\lambda| - m, m)}^{(2)} \otimes W_m^{(2)}.$$

Here $W_0, W_1, \dots$ are multiplicity spaces. The summands $V_{(|\lambda| - m, m)}^{(2)} \otimes W_m^{(2)}$ are distinguished by eigenvalues of the central element $I^{(2)}$:

$$(5.12) \quad I^{(2)} \Big|_{V_{(|\lambda| - m, m)}^{(2)} \otimes W_m^{(2)}} = |\lambda|^2 + |\lambda| + 2m(m - |\lambda| - 1).$$

Set

$$(V_{l_1}^{(2)} \otimes \dots \otimes V_{l_n}^{(2)})[m_1, m_2]_m = (V_{(|\lambda| - m, m)}^{(2)} \otimes W_m^{(2)}) \cap (V_{l_1}^{(2)} \otimes \dots \otimes V_{l_n}^{(2)})[m_1, m_2].$$

It follows from [TV1, Section 2.5] that

$$(5.13) \quad B_{12}^{(2)}(t) \Big|_{(V_{l_1}^{(2)} \otimes \dots \otimes V_{l_n}^{(2)})[m_1, m_2]_m} = \prod_{j=m}^{m_2-1} \frac{t + m_2 - j}{t - m_1 + j}.$$

Consider the decomposition of the $\mathfrak{gl}_n$ -module $V_{m_1}^{(n)} \otimes V_{m_2}^{(n)}$ into a direct sum of irreducibles:

$$(5.14) \quad V_{m_1}^{(n)} \otimes V_{m_2}^{(n)} = \bigoplus_{m=0}^{\min(m_1, m_2)} V_{(m_1+m_2-m, m, 0, \dots, 0)}^{(n)}.$$

The summands $V_{(m_1+m_2-m, m, 0, \dots, 0)}^{(n)}$ are distinguished by eigenvalues of the central element $I^{(n)}$:

$$(5.15) \quad I^{(n)} \Big|_{V_{(m_1+m_2-m, m, 0, \dots, 0)}^{(n)}} = \\ = (m_1 + m_2)^2 + (n-1)(m_1 + m_2) + 2m(m - m_1 - m_2 - 1).$$

Since the R -matrix $R^{(n)}(u)$ is $\mathfrak{gl}_n$ -invariant, see (4.1), it acts as a scalar on each summand of the decomposition (5.14). The eigenvalues of $R^{(n)}(t)$ can be found from (4.2), see [KRS], [KR]:

$$(5.16) \quad R^{(n)}(t) \Big|_{V_{(m_1+m_2-m, m, 0, \dots, 0)}^{(n)}} = \prod_{j=0}^{m-1} \frac{t - m_1 + j}{t + m_2 - j}.$$

Set

$$(V_{m_1}^{(n)} \otimes V_{m_2}^{(n)})[l_1, \dots, l_n]_m = V_{(m_1+m_2-m, m, 0, \dots, 0)}^{(n)} \cap (V_{m_1}^{(n)} \otimes V_{m_2}^{(n)})[l_1, \dots, l_n].$$

Let $\mu = (m_1, m_2)$. Lemma 5.7 and formulae (5.12), (5.15) imply that under isomorphisms (5.5) and (5.6) the images of $(V_{l_1}^{(2)} \otimes \dots \otimes V_{l_n}^{(2)})[m_1, m_2]_m$ and $(V_{m_1}^{(n)} \otimes V_{m_2}^{(n)})[l_1, \dots, l_n]_m$ in the space $P_{kn}[\lambda, \mu]$ coincide. Comparing now formulae (5.7), (5.13) and (5.16), we get the required statement. $\square$

References

- [B] P. Baumann, *q-Weyl group of a q-Schur algebra*, Preprint (1999).
- [CP] C. De Concini and C. Procesi, unpublished (1995).
- [EV] P. Etingof and A. Varchenko, *Dynamical Weyl groups and applications*, Adv. Math. **167** (2002), no. 1, 74–127.
- [FMTV] G. Felder, Ya. Markov, V. Tarasov and A. Varchenko, *Differential equations compatible with KZ equations*, Math. Phys., Analysis and Geometry **3** (2000), 139–177.
- [FR] I. B. Frenkel and N. Yu. Reshetikhin, *Quantum affine algebras and holonomic difference equations*, Comm. Math. Phys. **146** (1992), 1–60.
- [FV] G. Felder and A. Varchenko, *Three formulas for eigenfunctions of integrable Schrödinger operators*, Compositio Math. **107** (1997), 143–175.

- [H] R. Howe, *Perspectives on invariant theory: Schur duality, multiplicity-free actions and beyond*, Israel Math. Conf. Proc. **8** (1995), 1–182.
 - [Ha1] J. Harnad, *Dual isomonodromic deformations and moment maps to loop algebras*, Comm. Math. Phys. **166** (1994), no. 2, 337–366.
 - [Ha2] J. Harnad, *Quantum isomonodromic deformations and the Knizhnik-Zamolodchikov equations*, CRM Proc. Lecture Notes **9** (1996), 155–161.
 - [KR] P. P. Kulish and N. Yu. Reshetikhin, *On GL_3 -invariant solutions of the Yang-Baxter equation and associated quantum systems*, J. Soviet Math. **34** (1986), 1948–1971.
 - [KRS] P. P. Kulish, N. Yu. Reshetikhin and E. K. Sklyanin, *Yang-Baxter equation and representation theory I*, Lett. Math. Phys. **5** (1981), 393–403.
 - [MNO] A. Molev, M. Nazarov and G. Olshanski, *Yangians and classical Lie algebras*, Russian Math. Surveys **51** (1996), 205–282.
 - [MV] Ya. Markov and A. Varchenko, *Hypergeometric solutions of trigonometric KZ equations satisfy dynamical difference equations*, Adv. Math. **166** (2002), no. 1, 100–147.
 - [R] N. Reshetikhin, *The Knizhnik-Zamolodchikov system as a deformation of the isomonodromy problem*, Lett. Math. Phys. **26** (1992), no. 3, 167–177.
 - [ST] Yu. F. Smirnov and V. N. Tolstoy, *Extremal projectors for usual, super and quantum algebras and their use for solving Yang-Baxter problem*, in Selected topics in QFT and Mathematical Physics, (Proc. of the V-th Intern. Conf., Liblice, Czechoslovakia, June 25–30, 1989), World Scientific, Singapore, 1990.
 - [TL] V. Toledano-Laredo, *A Kohno-Drinfeld theorem for quantum Weyl groups*, Duke Math. J. **112**, no. 3, 421–451.
 - [TV1] V. Tarasov and A. Varchenko, *Difference equations compatible with trigonometric KZ differential equations*, Int. Math. Res. Notices (2000), no. 15, 801–829..
 - [TV2] V. Tarasov and A. Varchenko, *The dynamical differential equations compatible with the rational qKZ equations*, in preparation.
 - [Zh1] D. P. Zhelobenko, *Compact Lie groups and their representations*, Transl. Math. Mono., vol. 40, AMS, Providence RI, 1983.
 - [Zh2] D. P. Zhelobenko, *Extremal cocycles on Weyl groups*, Funk. Analiz i ego pril. **21** (1987), 183–192.
- D. P. Zhelobenko, *Extremal projectors and generalized Mickelsson algebras on reductive Lie algebras*, Math. USSR, Izvestiya **33** (1989), 85–100.